

LETTER TO THE EDITOR

Circulating Tumor Cells Serve as Potential Biomarkers for Hepatocellular Carcinoma Patient with Low Alpha-Fetoprotein

Yang Zhao^{2, #}, Shengtao Tian^{1, #}, Zhenhu Ma¹, Beibei Su¹, Hui Xie¹

These authors contributed equally to this work

¹ Department of Interventional Therapy, Fifth Medical Center of Chinese PLA General Hospital, Beijing, China

² Department of Cardiovascular Surgery, Special Medical Center for Strategic Support Forces, Beijing, China

(Clin. Lab. 2021;67:xx-xx. DOI: 10.7754/Clin.Lab.2021.210108)

Correspondence:

Hui Xie

Department of Interventional Therapy

Fifth Medical Center of

Chinese PLA General Hospital

100039 Beijing

China

Email: xh302jr@126.com

Supplementary Tables and Figures

Table S1. Patients' status post RFA (AFP elevated patients).

Patient information				AFP before RFA		AFP post RFA		Status post RFA	
Patient number	Diagnosis date	Gender	Age	Examination date	AFP (ng/mL)	Examination date	AFP (ng/mL)	Duration	Status
4	2020/3/11	male	52	2020/2/25	3.1	2020/3/24	5.19	6 months	CR
5	2020/4/7	male	71	2020/4/2	2.37	2020/4/8	2.39	3 months	CR
6	2020/4/7	male	67	2020/3/31	2.76	2020/4/9	3.15	5 months	CR
13	2019/12/17	male	71	2019/12/9	1.22	2020/5/21	1.31	5 months	CR
22	2020/3/19	male	81	2020/5/20	4.88	2020/7/1	6.88	4 months	CR




Figure S1. Epithelial and mixed types of CTC count alterations before or after RFA.

A - CTC counts (epithelial type) before and after RFA in HCC patients with less than 10 ng/mL AFP. B - CTC counts (mixed type) before and after RFA in HCC patients with less than 10 ng/mL AFP.